

Sequences and limits

A sequence is an infinite, ordered collection of real numbers

$$a_1, a_2, a_3, \dots, a_n, \dots$$

notation: or (doesn't have to start at $n=1$)

EXAMPLES

$$\begin{array}{ll} 5, 7, 9, 11, \dots & a_n = \\ 3, 6, 9, 12, \dots & a_n = \end{array}$$

$a_1 = \left\{ \begin{array}{l} \text{Arithmetic} \\ \text{Sequences} \end{array} \right.$

$$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \quad a_n = \quad \text{harmonic sequence}$$

$$-1, 1, -1, 1, \dots \quad b_n =$$

$$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots \quad a_n = \quad \text{geometric sequence}$$

$$1, 1, 2, 3, 5, 8, 13, \dots \quad \text{Fibonacci sequence}$$

$a_1 = a_2 = 1$

Limits


DEFINITION 8.2. (*Intuitive definition of the limit of a sequence*)

Let $\{a_n\}$ be a sequence and L be a real number. We say that $\{a_n\}$ has a limit L if we can make a_n arbitrarily close to L by taking n to be sufficiently large. We denote this situation by


$$\lim_{n \rightarrow \infty} a_n = L.$$

We say that $\{a_n\}$ is convergent if $\lim_{n \rightarrow \infty} a_n$ exists; otherwise we say that $\{a_n\}$ is divergent.


EXAMPLES



$$a_n = 1 + \frac{1}{n}$$



$$b_n = (-1)^n$$



$$a_n = 1 + \frac{(-1)^n}{n}$$

The formal definition:

$\lim_{n \rightarrow \infty} a_n = L$ if for all $\epsilon > 0$ there exists N such that if $n > N$

then $|a_n - L| < \epsilon$

THEOREM 8.4 (Limit laws). Let $\{a_n\}$ and $\{b_n\}$ be convergent sequences with $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$. Then:

1. $\lim_{n \rightarrow \infty} (a_n \pm b_n) = a \pm b$.
2. $\lim_{n \rightarrow \infty} (c a_n) = c a$ for any constant $c \in \mathbb{R}$.
3. $\lim_{n \rightarrow \infty} (a_n b_n) = a b$.
4. If $b \neq 0$ and $b_n \neq 0$, for all n then $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}$.

$$\text{Ex: } \lim_{n \rightarrow \infty} \frac{n^2 - n + 1}{3n^2 + 2n - 1} =$$

\approx

$$= \frac{1}{3}$$

THEOREM 8.5 (The squeeze theorem or the sandwich theorem). Let $\{a_n\}$, $\{b_n\}$ and $\{c_n\}$ be sequences such that $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = a$ and

$$a_n \leq b_n \leq c_n$$

for all $n \geq 1$. Then the sequence $\{b_n\}$ is also convergent and $\lim_{n \rightarrow \infty} b_n = a$.


$$\text{Ex: } \lim_{n \rightarrow \infty} \frac{\cos n}{n}$$

Diverging to infinity

We say a sequence a_n diverges to infinity ($\lim_{n \rightarrow \infty} a_n = \infty$ or $a_n \rightarrow \infty$)

if for any number $M > 0$ there is a term in the sequence after which all terms are bigger than M .

Formally: for all $M > 0$, there exists N such that if $n > N$ then $a_n > M$.



Similarly, $a_n \rightarrow -\infty$ if for all $M < 0$ there exist N such that if $n > N$ then $a_n < M$

Warning: don't treat ∞ like a number/actual limit

if $a_n \rightarrow \infty$ and $b_n \rightarrow -\infty$ it does NOT follow that $\lim_{n \rightarrow \infty} (a_n + b_n) = 0$

$$\text{eg: } a_n = n$$

$$b_n = -2n$$

$$a_n + b_n =$$

Some properties:

If $a_n \neq 0$ for all n , then $a_n \rightarrow 0 \Leftrightarrow \frac{1}{|a_n|} \rightarrow \infty$

If $a_n > 0$ then $a_n \rightarrow \infty \Leftrightarrow \frac{1}{a_n} \rightarrow 0$. (1)

Example the geometric sequence

$$\underline{a_n = r^n} \quad \text{if } r > 1 \quad (2)$$

$$r < -1$$

$$0 < r < 1$$

$$(2) \Rightarrow \Leftrightarrow (1)$$

The monotone sequences theorem

Bounded sequences

A sequence is called
bounded above if there is a number A such that $a_n \leq A$ for all n
bounded below if there is B s.t. $a_n \geq B$
bounded if it is bounded above and below

e.g.: $a_n = n$ bounded below by 1, not bounded above

$a_n = \frac{1}{n}$ bounded above by 1, bounded below by 0,
bounded

$a_n = n \sin n$ not bounded above
not bounded below.

Every convergent sequence is bounded, but the converse is not true

e.g.: $(-1)^n$ bounded but divergent

Upper (and lower bounds) are not unique

We call the smallest (least) upper bound the supremum $\sup\{a_n\}$

The greatest lower bound is called the infimum $\inf\{a_n\}$

If there is a maximum then $\sup = \max$, if there is a minimum
then $\inf = \min$

e.g.: $a_n = \frac{1}{n}$ has $\sup\{a_n\} = 1 = \max\{a_n\}$
 $\inf\{a_n\} = 0$, no minimum.

A sequence is called monotone if it is

non-decreasing: $a_1 \leq a_2 \leq a_3 \leq \dots a_n \leq a_{n+1} \dots$ or

non-increasing: $a_1 \geq a_2 \geq a_3 \geq \dots a_n \geq a_{n+1} \dots$

THEOREM 8.15 (The monotone sequences theorem). *If the sequence $\{a_n\}_{n=1}^{\infty}$ is non-decreasing and bounded above, then the sequence is convergent and*


$$\lim_{n \rightarrow \infty} a_n = \sup(\{a_n\}).$$

If $\{a_n\}_{n=1}^{\infty}$ is non-increasing and bounded below, then $\{a_n\}$ is convergent and

$$\lim_{n \rightarrow \infty} a_n = \inf(\{a_n\}).$$

That is, every monotone bounded sequence is convergent.

EXAMPLE



$$a_n = 1 + \frac{1}{n}$$

inf

$$\frac{1}{n+1} < \frac{1}{n}$$

$$1 + \frac{1}{n+1} < 1 + \frac{1}{n}$$

$$a_{n+1} < a_n$$

Infinite series

An infinite series is the sum of the terms in a sequence.

$$\sum_{n=1}^{\infty} a_n = a_1 + a_2 + \dots + a_n + \dots$$

For each $n \geq 1$, the n^{th} partial sum is

$$S_n = a_1 + a_2 + \dots + a_n = \sum_{i=1}^n a_i$$

The partial sums of an infinite series form a sequence:

$$S_1, S_2, S_3, \dots, S_n, \dots \quad (S_n)$$

If this sequence has a limit $\lim_{n \rightarrow \infty} S_n = S$, then we say the infinite series is convergent and write $\sum_{n=1}^{\infty} a_n = S$

i.e. $S = \lim_{n \rightarrow \infty} \sum_{i=1}^n a_i = \lim_{n \rightarrow \infty} (a_1 + a_2 + \dots + a_n)$

EXAMPLE $1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$

$$\xleftarrow{\hspace{1cm}} \quad | \quad \xrightarrow{\hspace{1cm}}$$

recall: geometric sequence $(1, r, r^2, \dots, r^n, \dots)$, $r \in \mathbb{R}$

geometric series

$$\sum_{i=0}^{\infty} r^i = 1 + r + r^2 + \dots$$

does this series converge?

the n^{th} partial sum is $S_n = 1 + r + r^2 + \dots + r^{n-1}$

$$\Rightarrow r S_n =$$

$$S_n - r S_n =$$

$$S_n(1 - r) =$$

$$S_n = \quad \text{for } r \neq 1$$

does the sequence s_n converge? This depends on r :

if $|r| < 1$ then $\lim_{n \rightarrow \infty} r^n = 0$ so

$$s_n = \frac{1 - r^n}{1 - r} \xrightarrow{n \rightarrow \infty} = \quad (\text{convergent})$$

if $r > 1$ then $r^n \xrightarrow{n \rightarrow \infty}$ (eg: $r = \frac{1}{2}$, $\frac{1}{1-\frac{1}{2}} = 2$)

and since $s_n = 1 + r + r^2 + \dots + r^n > r^n$

it follows that $s_n \rightarrow \infty$ too (divergent)

if $r < 1$ then $\lim_{n \rightarrow \infty} r^n$ doesn't exist so

$$s_n = \frac{1 - r^n}{1 - r} = \frac{1}{1 - r} - \frac{1}{1 - r} r^n \quad \text{also diverges}$$

if $r = 1$ then $s_n =$ divergent

if $r = -1$ then the sequence $(s_n) =$ which is divergent.

To summarise: the geometric series is convergent iff $|r| < 1$

THEOREM 8.27. If the infinite series $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are convergent,

then the series $\sum_{n=1}^{\infty} (a_n \pm b_n)$ and $\sum_{n=1}^{\infty} c a_n$ (for any constant $c \in \mathbb{R}$) are also convergent with

$$\sum_{n=1}^{\infty} (a_n \pm b_n) = \sum_{n=1}^{\infty} a_n \pm \sum_{n=1}^{\infty} b_n \quad \text{and} \quad \sum_{n=1}^{\infty} c a_n = c \sum_{n=1}^{\infty} a_n.$$

Example

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \left(\left(-\frac{2}{3}\right)^n + \frac{3}{4^{n+1}} \right) \\
 &= \sum_{n=0}^{\infty} \left(\left(-\frac{2}{3}\right)^n + \frac{3}{4} \left(\frac{1}{4}\right)^n \right) \quad \begin{matrix} \text{geometric series } \sum r^n \\ \rightarrow \frac{1}{1-r} \end{matrix} \\
 &= \sum_{n=0}^{\infty} \left(-\frac{2}{3}\right)^n + \frac{3}{4} \sum_{n=0}^{\infty} \left(\frac{1}{4}\right)^n
 \end{aligned}$$

$$= \frac{1}{1+\gamma_3} + \frac{3}{4} \cdot \frac{1}{1-\gamma_4}$$

$$= \frac{3}{5} + \frac{3}{4} \cdot \frac{4}{3} = \frac{8}{5}$$

Divergence test

A bit of logic: If P then Q \leftarrow suppose this is true

contrapositive: if not Q then not P \leftarrow then so is this

converse: if Q then P \leftarrow but this may not be

THEOREM 8.22. If the series $\sum_{n=1}^{\infty} a_n$ is convergent, then $\lim_{n \rightarrow \infty} a_n = 0$.

contrapositive: if $\lim_{n \rightarrow \infty} a_n \neq 0$ then the series is not convergent

TRUE "DIVERGENCE TEST"

converse: if $\lim_{n \rightarrow \infty} a_n = 0$ then $\sum_{n=1}^{\infty} a_n$ is convergent

FALSE

Example Is $\sum_{n=1}^{\infty} \frac{n^2-1}{2n^2+2}$ convergent?

$$\lim_{n \rightarrow \infty} \frac{n^2-1}{2n^2+2} = \lim_{n \rightarrow \infty} \frac{1 - \frac{1}{n^2}}{2 + \frac{2}{n^2}} = \frac{1}{2}$$

therefore divergent by divergence test

Is $\sum_{n=1}^{\infty} \frac{1}{n}$ (harmonic series) convergent?

$\lim_{n \rightarrow \infty} \frac{1}{n} = 0$ can't decide based only on the divergence test.

(1350 Oresme)

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \dots$$

$$> 1 + \frac{1}{2} +$$

$$= 1 + \frac{1}{2} + \text{divergent}$$